



Characteristics of Nest Cavities and Nest Trees of the Red-Breasted Sapsucker in Coastal Montane Forests (Características de la Cavidad y el Árbol Utilizado para Anidar por Parte de *Sphyrapicus ruber* en Bosques Montanos Costaneros)

Author(s): Jeffrey B. Joy

Source: *Journal of Field Ornithology*, Vol. 71, No. 3 (Summer, 2000), pp. 525-530

Published by: Blackwell Publishing on behalf of Association of Field Ornithologists

Stable URL: <http://www.jstor.org/stable/4514517>

Accessed: 02/12/2008 14:52

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=afo>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We work with the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact support@jstor.org.



Association of Field Ornithologists and Blackwell Publishing are collaborating with JSTOR to digitize, preserve and extend access to *Journal of Field Ornithology*.

CHARACTERISTICS OF NEST CAVITIES AND NEST TREES OF THE RED-BREASTED SAPSUCKER IN COASTAL MONTANE FORESTS

JEFFREY B. JOY

Mt. Cain Montane Research Station
598 Millstream Lake Road
Victoria, British Columbia
V9E 1E7, Canada

Abstract.—From May through July 1997 and 1998, I quantified nest tree and cavity characteristics of 32 Red-breasted Sapsucker (*Sphyrapicus ruber*) nests in the high-elevation coastal forests of northern Vancouver Island, British Columbia. Nests were located in western white pine (*Pinus monticola*), hemlock (*Tsuga sp.*), and Douglas-fir (*Pseudotsuga menziesii*) snags. Nest trees were significantly taller and had a greater DBH than random snags. Nest height was positively correlated with tree height. All nests were in dead trees. The orientation of nest cavity entrances did not differ from random orientations. When choosing nest sites, Red-breasted Sapsuckers are likely balancing predation risk (decreasing at higher nest heights) and adequate nest space and insulation (greater with increasing diameter at lower nest heights).

CARACTERÍSTICAS DE LA CAVIDAD Y EL ÁRBOL UTILIZADO PARA ANIDAR POR PARTE DE *SPHYRAPICUS RUBER* EN BOSQUES MONTANOS COSTANEROS

Sinopsis.—De mayo a julio de 1997 y 1998, cuantifiqué las características de la cavidad y el árbol utilizado para anidar por parte de 32 parejas de *Sphyrapicus ruber* en bosques montanos costaneros de Vancouver, Columbia Británica. Los nidos fueron localizados en árboles de *Pinus monticola*, *Tsuga sp.* y troncos partidos de *Pseudotsuga menziesii*. Los árboles utilizados para anidar resultaron significativamente más altos y con mayor DBH que los troncos partidos tomados al azar. La altura del nido estuvo positivamente correlacionada con la altura del árbol. Todos los nidos fueron encontrados en vegetación muerta. La orientación de la entrada de la cavidad del nido no resultó diferente a orientaciones al azar. Cuando las aves están seleccionando el lugar para anidar parecen poner en balance el riesgo de depredación (disminuye con la altura a que se construye el nido) con la insulación y el espacio adecuado para la construcción del nido (mayor con el incremento en el diámetro del tronco a menor altura).

The Red-breasted Sapsucker (*Sphyrapicus ruber*) is the most common primary cavity excavator in coastal montane forests of the Pacific Northwest (Bryant 1995). Because sapsucker excavations can provide important nesting and feeding opportunities for other bird species (Daily 1993, Walters 1996), the Red-breasted Sapsucker may be a critical component of coastal montane forest bird communities. Unfortunately, little is known about how timber harvesting, which reduces availability of suitable nest trees (Zarnowitz and Manuwal 1985, Martin and Eadie 1999), might impact this important species.

Because timber removal at higher elevations in coastal British Columbia is increasing (Whitehead 1995), a study of nest site selection by Red-breasted Sapsuckers in this region was needed. Therefore, the specific objectives of my study were to quantify Red-breasted Sapsucker nest-tree and cavity-site dimensions in coastal montane forests.










Live		Dead					Dead Fallen	
1	2	3	4	5	6	7	8	9
								
Live/healthy; No decay	Live/unhealthy; internal decay or growth deficiencies (including insect damage, broken tops); dying tree	Dead; needles or twigs may be present; roots sound	Dead; no needles/twigs; 50% of branches lost; loose bark; top usually broken; roots stable	Dead; most branches/bark absent; some internal decay; roots of larger trees stable	Dead; no branches or bark; uppermost heartwood absent; some internal decay; upper bark decay more advanced; internal roots of larger trees softening; smaller trees unstable	Dead; extensive internal decay; outer shell may be hard; internal roots completely disintegrated; hollow or nearly hollow shells		Detritic; downed trees or snags

FIGURE 1. Wildlife tree classification system used to describe nest and non-nest trees. Numbers 1 through 9 denote decay classes the characteristics of which are pictured and described beneath.

METHODS

This study took place on Mt. Cain and Maquilla Peak (50°13'N, 126°18'W), Vancouver Island, British Columbia. The study area encompassed 11,800 ha of coastal montane forest. The landscape consists of managed and unmanaged stands of coastal montane forest, 600–1650 m in elevation. The dominant tree species included mountain hemlock (*Tsuga mertensiana*), western hemlock (*T. heterophylla*), yellow cedar (*Chamaecyparis nootkatensis*), and amabilis fir (*Abies amabilis*).

I searched for active Red-breasted Sapsucker nests between May and July, 1997 and 1998. Nests were considered active when nestlings could be heard. If adult birds were observed entering a cavity it was revisited until nestlings were heard or until it was clearly not an active nest. After the young had fledged in late July and early August, I quantified characteristics of the active nest trees and nest cavities. I recorded tree species based on remnant bark, wood condition, and tree form. Internal tree decay was evaluated from increment bore samples of heartwood and sapwood. Wildlife tree class from 1 (live, healthy) to 9 (very rotten) was assigned based on tree decay characteristics (Guy and Manning 1994, Fig. 1). Tree height was measured using a clinometer and tape measure.

I reached cavity nests using nearby live trees and climbing equipment. From each active nest I obtained the following data: entrance vertical and horizontal diameter, cavity depth, cavity orientation, cavity height, and diameter at cavity height. Cavity depth was measured using a plumb bob lowered into the cavity.

Comparative data were gathered from 135 10 × 10-m plots, systematically distributed at 300-m intervals throughout unmanaged forests in the study area. At each plot, I obtained the following data for all trees >2 cm dbh: tree species, DBH, tree height, bark retention characteristics, and stage of decay. For comparative purposes random samples of trees >17 cm DBH were drawn from these data since Crockett and Hadow (1975) found that this is the minimum diameter that could provide adequate space for a nesting sapsucker.




FIGURE 2. Diameter classes of trees used for nesting by Red-breasted Sapsuckers and of trees available in the Mt. Cain and Maquilla Peak study area of British Columbia, 1997 and 1998. Mean dbh of nest trees ($\bar{x} = 93.3 \text{ cm} \pm 4.6 \text{ SE}$, $n = 31$) was significantly greater than non-nest trees ($\bar{x} = 32.02 \text{ cm} \pm 2.2 \text{ SE}$, $n = 98$).

RESULTS

I located 32 active Red-breasted Sapsucker nests in 31 snags (17 in 1997 and 15 in 1998). No nest cavities were reused in the second year. Nests were located in western white pine (*Pinus monticola*) ($n = 7$), hemlock ($n = 6$), Douglas fir (*Pseudotsuga menziesii*) ($n = 1$), and unidentified snags ($n = 17$). Mean DBH of nest trees was significantly greater than that of non-nest trees $>17 \text{ cm dbh}$ ($t = 10.54$, $df = 125$, $P < 0.001$, Fig. 2.). Mean nest tree height ($\bar{x} = 32.5 \text{ m} \pm 2.4 \text{ SE}$, $n = 31$, range 16.8–47.1) was also significantly greater than the mean height of random trees ($\bar{x} = 14.75 \text{ m} \pm 0.81 \text{ SE}$, $n = 88$, $t = 10.45$, $df = 118$, $P < 0.001$). The distribution of wildlife tree classes differed significantly from expected ($G = 23.95$, $df = 6$, $P < 0.05$, Fig. 3). Sapsuckers selected only dead trees. I measured the cavity entrance dimensions and depth of 22 of the 32 active sapsucker nests (Table 1). Nest cavity orientation did not differ significantly from a random distribution (Rayleigh's test; $R = 9.752$, $P < 0.05$, $n = 32$). Nest height averaged 17.2 m and was positively correlated with tree height ($r = 0.55$, $df = 30$, $P < 0.001$).

DISCUSSION

In the coastal montane forests of Northern Vancouver Island, Red-breasted Sapsuckers consistently chose nest trees of larger diameter than non-nest trees, as has been found with other sapsuckers (Dobkin et al. 1995) and other small woodpeckers (Lundquist and Mariani 1991). The choice of large diameter nest trees may reflect an attempt by this species to maximize nest space, thermal insulation, and/or protection from predation. Red-breasted Sapsuckers require large nest cavities to accommodate large clutches and up to six nestlings (Campbell et al. 1990). The




FIGURE 3. Wildlife tree classes of trees used for nesting by the Red-breasted Sapsucker and of a random selection of non-nest trees in the Mt. Cain and Maquilla Peak study area of British Columbia, 1997 and 1998.

greater insulation provided by larger snags may be beneficial to cavity nesting birds in high elevation forests where night temperatures can be below freezing well into the breeding season (pers. obs.). Additionally, large diameter nest trees may prohibit predators from reaching the nest by allowing cavities to have thicker walls (cf. Kilham 1971, Harestad and Keisker 1989). The preference for tall trees and high nest placement exhibited by Red-breasted Sapsuckers and other sapsuckers (Dobkin et al. 1995) likely reflects an adaptation to avoid predation by terrestrial predators. Rates of predation and nest failure decrease among cavity nesters with increasing nest height (Nilsson 1984, Li and Martin 1991). Overall, Red-breasted Sapsuckers likely choose nest sites that balance predation risk (which decreases at greater nest heights) and the availability of adequate nest space and insulation (greater at lower nest heights). Red-breasted Sapsuckers chose to nest in dead trees with internal decay. Decayed heartwood is essential for most primary cavity nesting birds to excavate nest holes (Harestad and Keisker 1989).

As mature forests are logged, the population sizes of the cavity nesting

TABLE 1. Means, standard errors, and ranges of nest-cavity attributes of Red-breasted Sapsuckers from the coastal montane forests of northern Vancouver Island, British Columbia, Canada.

Nest tree or cavity variable	n	\bar{x}	SE	Range
Diameter at cavity height (cm)	22	75.0	5.0	32.9–124.8
Cavity vertical diameter (cm)	22	4.7	0.1	4.3–5.4
Cavity horizontal diameter (cm)	22	4.6	0.1	3.9–5.7
Cavity depth (cm)	22	24.8	9.6	14.1–34.2
Cavity height (m)	32	17.2	1.5	5.8–31.7

species that inhabit them are reduced, particularly those species which prefer large trees for nesting such as the Red-breasted Sapsucker. It is therefore important that snag retention is used to create some suitable habitat for such species in regenerating forests. It is now clear that primary cavity nesting species serve as a keystone group in northern forests (Daily et al. 1993, Walters 1996). My data suggest that in designing wildlife tree patches foresters should include tall large diameter snags and nearly dead trees to ensure suitable nest sites are available for Red-breasted Sapsuckers.

ACKNOWLEDGMENTS

I thank C. R. Chandler, D. J. Ingold, J. N. M. Smith, and R. van den Driessche for comments which greatly improved this manuscript and Harry van Oort for assistance in collecting the field data. Thanks also to the British Columbia Wildlife Tree Committee for Figure 1. I am grateful to the British Columbia Ministry of Forests, Research Branch and British Columbia Ministry of Environment, Lands, and Parks, Region One, and Forest Renewal British Columbia for financial support. R. S. McNay provided support throughout the study. Canadian Forest Products Ltd. also provided support.

LITERATURE CITED

- BRYANT, A. 1995. Putting MASS in Context: bird communities in Vancouver Island old-growth forests. Pp. 101–112, in J. T. Arnott, W. J. Beese, A. K. Mitchell, and J. Peterson, eds. *Montane Alternative Silvicultural Systems (MASS): proceedings of a workshop*, Courtney, British Columbia, 7–8 June 1995. Canadian Forest Service, FRDA Report No. 238.
- CAMPBELL, R. W., N. K. DAWE, I. McTAGGART-COWAN, J. M. COOPER, G. W. KAISER, AND M. C. E. McNALL. 1990. *The Birds of British Columbia*, Vol. II. Royal British Columbia Museum, Victoria, British Columbia.
- CROCKETT, A. B., AND H. H. HADOW. 1975. Nest site selection by Williamson and Red-naped Sapsuckers. *Condor* 77:365–368.
- DAILY, G. C. 1993. Heartwood decay and vertical distribution of Red-naped Sapsucker nest cavities. *Wilson Bull.* 105:674–679.
- , P. R. EHRLICH, AND N. M. HADDAD. 1993. Double keystone bird in a keystone species complex. *Proc. Natl. Acad. Sci.* 90:592–594.
- DOBKIN, D. S., A. C. RICH, J. A. PRETARE, AND W. H. PYLE. 1995. Nest-site relationships among birds of riparian and snowpocket aspen woodlands in the northwestern Great Basin. *Condor* 97:694–707.
- GUY, S., AND T. MANNING. 1995. *Wildlife/danger tree assessor's course workbook*. Canada-British Columbia Partnership Agreement on Forest Resource Development. Ministry of Forests. Victoria, British Columbia. 158 pp.
- HARESTAD, A. S., AND D. G. KEISKER. 1989. Nest tree use by primary cavity-nesting birds in south central British Columbia. *Can. J. of Zool.* 67:1067–1073.
- KILHAM, L. 1971. Reproductive behavior of Yellow-bellied Sapsuckers. I. Preference for nesting in *Fomes* infected aspens and nest hole interrelations with flying squirrels, raccoons, and other animals. *Wilson Bull.* 83:159–171.
- LI, P., AND T. E. MARTIN. 1991. Nest-site selection and nesting success of cavity-nesting birds in high elevation forest drainages. *Auk* 108:405–418.
- LUNDQUIST, R. W., AND J. M. MARIANI. 1991. Nesting habitat and abundance of snag dependent birds in the Southern Washington Cascade Range. Pp. 221–239, in K. B. Aubry, M. H. Brookes, J. K. Agee, J. F. Franklin, B. R. Noon, M. G. Raphael, R. M. Storm, and J. Verner, eds. *Wildlife and Vegetation of unmanaged Douglas Fir Forests*. Department of Agriculture Forest Service General Technical Report PNW-GTR-285. Pacific Northwest Research Station, Portland, Oregon.
- MARTIN, K. AND J. EADIE. 1999. Nest webs: A community-wide approach to the management and conservation of cavity-nesting forest birds. *For. Ecol. and Manage.* 115:243–257.

- NILSSON, S. G. 1984. The evolution of nest-site selection among hole-nesting birds: the importance of nest predation and competition. *Ornis Scandinavica* 15:165–175.
- WALTERS, E. 1996. Habitat and space use of the Red-naped Sapsucker, *Sphyrapicus nuchalis*, in the Hat Creek Valley, south-central British Columbia. M. Sc. Thesis, Univ. Victoria, Victoria, British Columbia.
- WHITEHEAD, R. 1995. The Montane Alternative Silvicultural Systems Project: A research and operations partnership. Pp. 1–2, in J. T. Arnott, W. J. Beese, A. K. Mitchell, and J. Peterson, eds. *Montane Alternative Silvicultural Systems (MASS): proceedings of a workshop*, Courtney, British Columbia, 7–8 June 1995. Canadian Forest Service, FRDA Report No. 238.
- ZARNOWITZ, J. AND D. MANUWAL. 1985. The effects of forest management on cavity-nesting birds in Northwestern Washington. *J. Wildl. Manage.* 49:255–263.

Received 18 Feb. 1999; accepted 8 Jul. 1999.